




The ‘Provenance Issue’: Challenges and Opportunities for Ecological Restoration

Dr Maurizio Rossetto




Threats: Habitat Degradation



Vegetation Clearing

Threats: Climate Change



Responses to Change

- ADAPT
 - Population size and the diversity within it sufficient for evolutionary response
 - Persist through plastic resilience
 - Multiple provenances with different adaptation
- MOVE
 - To newly available or remaining habitat
 - If capable and competitive
 - Can gene flow support adaptation
- LOCAL EXTINCTION



Responses to Change

- ADAPT
 - Population size and the diversity within it sufficient for evolutionary response
 - Persist through plastic resilience
 - Multiple provenances with different adaptive potential
- MOVE
 - To newly available or remaining habitat
 - If capable and competitive
 - Can gene flow support adaptation
- LOCAL EXTINCTION



Responses to Change


- ADAPT
 - Population size and the diversity within it sufficient for evolutionary response
 - Persist through plastic resilience
 - Multiple provenances with different adaptation
- MOVE
 - To newly available or remaining habitat
 - If capable and competitive
 - Can gene flow support adaptation
- LOCAL EXTINCTION



Why is the evolutionary context important?

- Short-term: genomic diversity maximises fitness
 - loss of diversity / heterozygosity causes inbreeding
 - increased frequency of deleterious genes causes inbreeding depression


Pigmy hippopotamus



Why is the evolutionary context important?

- Long-term: genomic diversity maximises adaptation potential
 - reduces vulnerability to environmental changes
 - increases potential for evolutionary differentiation (i.e. speciation)
 - outbreeding depression: loss of fitness resulting from mixing individuals from different provenances

Rutidosia leptorrhyncoides



Telopea speciosissima



Ecological Restoration




Aims to recover the structure, dynamics and evolutionary potential of an ecosystem

**MORE THAN
GARDENING!**



Ecological Restoration: Limitations


- **Sustainability:** unsuitable plantings are costly and demand high maintenance
- **Research:** limited evidence-based studies
- **Information Resources:** no central source




Can we Generalise? Assemblage

Sydney Sandstone Flora


Isopogon anemonifolius



Petrophile pulchella





Leptospermum trinervium






Can we Generalise? Taxonomy

11 species of *Elaeocarpus* in AWT




E. elliffii






E. foveolatus





E. bancroftii





E. largiflorens




E. carolinae




Can we Generalise? Taxonomy




Can we Generalise? Biogeographic Barriers




What about Temporal Variation?



September



October




November

Short flowering season needs ~10 days of 20°C



What about Temporal Variation?


Glacial cycles and *Telopea*



Telopea speciosissima




Telopea mungaensis




Telopea oreades



What about Temporal Variation?



Podocarpus elatus




What about Natural Admixture?



Permeable species boundaries in *Lomatia*




Lomatia myricoides




Hybrid



Lomatia silaifolia




What about Natural Admixture?




- 5 species of *Lomatia*: genetic diversity is distributed geographically rather than taxonomically

What about Natural Admixture?


- Polar bears



Local Adaptation vs. Drift: the Green Ashes



Local Adaptation vs. Drift: the Green Ashes



Differences between treatments significant P<0.05




Upper Blue Mountains

E. dendromorpha Mt Banks

Blue Mountains

Sydney and southern populations



E. cunninghamii
Upper Blue Mountains

Upper Blue Mountains

Based on 130 000 DArTseq presence/absence markers

Provenance: Challenges and Opportunities

- Defining provenances is a mechanism for quantifying biodiversity
- They are a complex mix of drift and adaptation, often resulting in temporal flux of non-equilibrium lineages and assemblages
- Preserving them within changing climate needs understanding of the driving processes



Provenance: Challenges and Opportunities

- Biodiversity preservation is about managing mostly unknown change
- Restoring 'past composition' is difficult and often ecologically unrealistic
- Ecological restoration should focus on re-establishing 'evolutionary resilient landscapes'



Provenance: Challenges and Opportunities

Conclusion

Provenances should not be used to define hard boundaries but to develop decision-making frameworks that are evolutionary relevant



Provenance: Challenges and Opportunities

Evolutionary-relevant decision frameworks can lead to:

- Preservation of unique local adaptations
- Assisted re-colonisation of suitable habitats
- Genetic rescue and / or targeted admixture

Florida Panther





**Restore
&
Renew**



The Royal
BOTANIC GARDENS
& *Domain Trust*

Restore & Renew: introduction

- **Community website:** definitive and readily accessible tool
- **Location-specific & generalisations:** operational support and future proofing
- **Globally pioneering:** world-first initiative approach, scale and scope



How does it work?

Species selection

200+ widely used species through extensive community consultation

Environmental modelling and sampling


Trained citizen science teams to support collection from 5,000+ sites

Evolutionary and ecological analyses


Next Generation DNA Sequencing will reveal unparalleled information on provenance, diversity and dynamics

Easy access: website

User-friendly guidelines empowering communities to deliver successful and self-sustainable restoration



Example: NSW Waratah




Waratah (*Telopea speciosissima*)

Three distinct provenances:

- Upland
- Coastal
- Southern

(plus areas of admixture)



Species selection:

NSW Waratah (*T. speciosissima*)


Site selection (*Lat-Long*):

-33.42S; 150.19E

Supporting information:

- Upland (green) provenance ([more](#)).
- Preferential outcrosser, so must maintain heterozygosity ([more](#)).
- To achieve this ideally sample from multiple sites within the provenance and multiple individuals within each site ([more](#)).
- For 'future proofing' click [here](#).

Shifts in suitable habitat for each gene type by 2050



Species selection:

NSW Waratah (*T. speciosissima*)

Site selection (*Lat-Long*):

-33.42S; 150.19E

Supporting information:

- Suitable habitat will shift with coastal gaining ([more](#)).
- Predicted natural movement is limited ([more](#)).
- At selected site upland and coastal provenances could be mixed ([more](#)).
- To maintain upland provenance need conservation introduction in new area ([more](#)).



Get Involved!

- Become a project partner/sponsor
- Host a collection group
- restore.renew@rbgsyd.nsw.gov.au

**Restore
&
Renew**



The Royal
BOTANIC GARDENS
& Domain Trust

Postdocs:

*H McPherson
M van der Merwe*

Postgrads:

*R Mellick
M Milner
E McIntosh
S Rutherford*

External Collaborators:

*D Crayn (JCU)
A Ford (CSIRO)
R Henry (UQ)
A Lowe (Uni. of Adel.)*

RBG&DT Collaborators:

*C Allen
R Kooyman
C Porter
J Siow*



Funding:

*ABRS, ARC, Bioplatforms, Friends of the Gardens,
Hermon Slade Foundation*

